

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:ssptacrs1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	MAR 31	IFICDB, IFIPAT, and IFIUIDB enhanced with new custom IPC display formats
NEWS	3	MAR 31	CAS REGISTRY enhanced with additional experimental spectra
NEWS	4	MAR 31	CA/CAPplus and CASREACT patent number format for U.S. applications updated
NEWS	5	MAR 31	LPCI now available as a replacement to LDPCI
NEWS	6	MAR 31	EMBASE, EMBAL, and LEMBASE reloaded with enhancements
NEWS	7	APR 04	STN AnaVist, Version 1, to be discontinued
NEWS	8	APR 15	WPIDS, WPINDEX, and WPIX enhanced with new predefined hit display formats
NEWS	9	APR 28	EMBASE Controlled Term thesaurus enhanced
NEWS	10	APR 28	IMSRESEARCH reloaded with enhancements
NEWS	11	MAY 30	INPAFAMDB now available on STN for patent family searching
NEWS	12	MAY 30	DGENE, PCTGEN, and USGENE enhanced with new homology sequence search option
NEWS	13	JUN 06	EPFULL enhanced with 260,000 English abstracts
NEWS	14	JUN 06	KOREAPAT updated with 41,000 documents
NEWS	15	JUN 13	USPATFULL and USPAT2 updated with 11-character patent numbers for U.S. applications
NEWS	16	JUN 19	CAS REGISTRY includes selected substances from web-based collections
NEWS	17	JUN 25	CA/CAPplus and USPAT databases updated with IPC reclassification data
NEWS	18	JUN 30	AEROSPACE enhanced with more than 1 million U.S. patent records
NEWS	19	JUN 30	EMBASE, EMBAL, and LEMBASE updated with additional options to display authors and affiliated organizations
NEWS	20	JUN 30	STN on the Web enhanced with new STN AnaVist Assistant and BLAST plug-in
NEWS	21	JUN 30	STN AnaVist enhanced with database content from EPFULL
NEWS	22	JUL 28	CA/CAPplus patent coverage enhanced
NEWS	23	JUL 28	EPFULL enhanced with additional legal status information from the epoline Register
NEWS	24	JUL 28	IFICDB, IFIPAT, and IFIUIDB reloaded with enhancements
NEWS	25	JUL 28	STN Viewer performance improved
NEWS	26	AUG 01	INPADOCDB and INPAFAMDB coverage enhanced
NEWS	27	AUG 13	CA/CAPplus enhanced with printed Chemical Abstracts page images from 1967-1998
NEWS	28	AUG 15	CAOLD to be discontinued on December 31, 2008
NEWS	29	AUG 15	CAPplus currency for Korean patents enhanced

NEWS EXPRESS JUNE 27 08 CURRENT WINDOWS VERSION IS V8.3,

AND CURRENT DISCOVER FILE IS DATED 23 JUNE 2008.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 11:17:09 ON 18 AUG 2008

=> file registry

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'REGISTRY' ENTERED AT 11:17:24 ON 18 AUG 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 17 AUG 2008 HIGHEST RN 1041629-70-2

DICTIONARY FILE UPDATES: 17 AUG 2008 HIGHEST RN 1041629-70-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH July 5, 2008.


Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>

Uploading C:\Program Files\STNEXP\Queries\10552065.str



```

chain nodes :
13 25 32 33
ring nodes :
1 2 3 4 5 6 7 8 9 10 11 12 14 15 16 17 18 19 20 21 22 23 24
26 27 28 29 30 31
chain bonds :
3-13 6-7 10-16 14-32 15-22 23-25 26-32 29-33
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12 14-15 14-18
15-16 16-17 17-18 19-20 19-24 20-21 21-22 22-23 23-24 26-27 26-31 27-28
28-29 29-30 30-31
exact/norm bonds :
10-16 14-15 14-18 15-16 16-17 17-18 26-27 26-31 26-32 27-28 28-29 29-30
30-31
exact bonds :
3-13 6-7 14-32 15-22 23-25 29-33
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12 19-20 19-24
20-21 21-22 22-23 23-24

```

```

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:CLASS 14:Atom 15:Atom 16:Atom 17:Atom 18:Atom 19:Atom
20:Atom 21:Atom 22:Atom 23:Atom 24:Atom 25:CLASS 26:Atom 27:Atom 28:Atom
29:Atom 30:Atom 31:Atom 32:CLASS 33:CLASS

```

L1 STRUCTURE UPLOADED

```

=> s l1 fam ful
FULL SEARCH INITIATED 11:17:45 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 89 TO ITERATE

```

```

100.0% PROCESSED      89 ITERATIONS                      1 ANSWERS
SEARCH TIME: 00.00.01

```

L2 1 SEA FAM FUL L1

=> file caplus
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
70.11	70.32

FULL ESTIMATED COST

FILE 'CAPLUS' ENTERED AT 11:17:51 ON 18 AUG 2008
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 18 Aug 2008 VOL 149 ISS 8
FILE LAST UPDATED: 17 Aug 2008 (20080817/ED)

Caplus now includes complete International Patent Classification (IPC) reclassification data for the second quarter of 2008.

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/legal/infopolicy.html>

=> s 12
L3 1 L2

=> d 13 ibib abs

L3 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2004:841775 CAPLUS
DOCUMENT NUMBER: 141:350163
TITLE: Preparation of arylpyrazoles as serotonin 5-HT2A and 5-HT2C receptor antagonists
INVENTOR(S): Schiemann, Kai; Ackermann, Karl-August; Arlt, Michael; Finsinger, Dirk; Schadt, Oliver; Van Amsterdam, Christoph; Bartoszyk, Gerd; Seyfried, Christoph
PATENT ASSIGNEE(S): Merck Patent GmbH, Germany
SOURCE: Ger. Offen., 102 pp.
CODEN: GWXXBX
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
DE 10315572	A1	20041014	DE 2003-10315572	20030405
AU 2004228120	A1	20041021	AU 2004-228120	20040308
CA 2521201	A1	20041021	CA 2004-2521201	20040308
WO 2004089931	A1	20041021	WO 2004-EP2353	20040308
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,				

GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
 LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI,
 NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,
 TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,
 BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE,
 ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI,
 SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
 TD, TG

EP 1626967 A1 20060222 EP 2004-718277 20040308
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK
 BR 2004009164 A 20060411 BR 2004-9164 20040308
 CN 1768051 A 20060503 CN 2004-80008572 20040308
 JP 2006522035 T 20060928 JP 2006-504584 20040308
 US 20060264419 A1 20061123 US 2005-552065 20051005
 PRIORITY APPLN. INFO.: DE 2003-10315572 A 20030405
 WO 2004-EP2353 W 20040308
 OTHER SOURCE(S): MARPAT 141:350163
 GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Preparation of title compds. I [X = CH, N; R1 = H, halo, (CH2)nHet, etc.; R2 = (CH2)nHet, (CH2)nAr, cycloalkyl, etc.; R3, R4 = H, (CH2)nCOHet, CHO, etc.; n = 0-5; Ar = (un)substituted Ph; Het = (un)substituted monoarom., bicyclic-heterocycle] and their pharmaceutically acceptable salts were prepared For example, sodium triacetoxyborohydride mediated reductive amination of 1-methyl-piperazine and aldehyde II, e.g., prepared from 2-fluoro- α,γ -dioxo-benzenebutanoic Et ester in 4-steps, afforded the dihydrochloride salt of arylpyrazole III. In 5-HT2A receptor binding assays, 167-examples of compds. I exhibited IC50 values ranging from 0.015-4.7x10⁻⁷M. Compds. I are claimed suitable as ligands of 5-HT receptors.

=> file registry
 COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
3.39	73.71

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-0.80	-0.80

CA SUBSCRIBER PRICE

FILE 'REGISTRY' ENTERED AT 11:18:26 ON 18 AUG 2008
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2008 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
 provided by InfoChem.

STRUCTURE FILE UPDATES: 17 AUG 2008 HIGHEST RN 1041629-70-2
 DICTIONARY FILE UPDATES: 17 AUG 2008 HIGHEST RN 1041629-70-2

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH July 5, 2008.

Please note that search-term pricing does apply when conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

=> s l1 sss ful

FULL SEARCH INITIATED 11:18:30 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 163 TO ITERATE

100.0% PROCESSED 163 ITERATIONS

4 ANSWERS

SEARCH TIME: 00.00.01

L4 4 SEA SSS FUL L1

=> file caplus

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

178.36

252.07

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

0.00

-0.80

FILE 'CAPLUS' ENTERED AT 11:18:39 ON 18 AUG 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 18 Aug 2008 VOL 149 ISS 8

FILE LAST UPDATED: 17 Aug 2008 (20080817/ED)

Caplus now includes complete International Patent Classification (IPC) reclassification data for the second quarter of 2008.

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/legal/infopolicy.html>

=> s l4

L5 2 L4

=> d l5 ibib abs 1-2

L5 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2004:841775 CAPLUS
 DOCUMENT NUMBER: 141:350163
 TITLE: Preparation of arylpyrazoles as serotonin 5-HT_{2A} and 5-HT_{2C} receptor antagonists
 INVENTOR(S): Schiemann, Kai; Ackermann, Karl-August; Arlt, Michael; Finsinger, Dirk; Schadt, Oliver; Van Amsterdam, Christoph; Bartoszyk, Gerd; Seyfried, Christoph
 PATENT ASSIGNEE(S): Merck Patent GmbH, Germany
 SOURCE: Ger. Offen., 102 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10315572	A1	20041014	DE 2003-10315572	20030405
AU 2004228120	A1	20041021	AU 2004-228120	20040308
CA 2521201	A1	20041021	CA 2004-2521201	20040308
WO 2004089931	A1	20041021	WO 2004-EP2353	20040308
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1626967	A1	20060222	EP 2004-718277	20040308
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
BR 2004009164	A	20060411	BR 2004-9164	20040308
CN 1768051	A	20060503	CN 2004-80008572	20040308
JP 2006522035	T	20060928	JP 2006-504584	20040308
US 20060264419	A1	20061123	US 2005-552065	20051005
PRIORITY APPLN. INFO.:			DE 2003-10315572	A 20030405
			WO 2004-EP2353	W 20040308
OTHER SOURCE(S): MARPAT 141:350163				
GI				

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Preparation of title compds. I [X = CH, N; R₁ = H, halo, (CH₂)_nHet, etc.; R₂ = (CH₂)_nHet, (CH₂)_nAr, cycloalkyl, etc.; R₃, R₄ = H, (CH₂)_nCOHet, CHO, etc.; n = 0-5; Ar = (un)substituted Ph; Het = (un)substituted monoarom., bicyclic-heterocycle] and their pharmaceutically acceptable salts were prepared For example, sodium triacetoxyborohydride mediated reductive amination of 1-methyl-piperazine and aldehyde II, e.g., prepared from 2-fluoro- α,γ -dioxo-benzenebutanoic Et ester in 4-steps, afforded the dihydrochloride salt of arylpyrazole III. In 5-HT_{2A} receptor binding assays, 167-examples of compds. I exhibited IC₅₀ values ranging from 0.015-4.7x10⁻⁷M. Compds. I are claimed suitable as ligands of 5-HT receptors.

ACCESSION NUMBER: 2004:841772 CAPLUS
 DOCUMENT NUMBER: 141:332186
 TITLE: Preparation of arylpyrazoles as serotonin 5-HT_{2A} and/or 5-HT_{2C} receptor antagonists.
 INVENTOR(S): Schadt, Oliver; Arlt, Michael; Finsinger, Dirk; Schiemann, Kai; Van Amsterdam, Christoph; Bartoszyk, Gerd; Seyfried, Christoph
 PATENT ASSIGNEE(S): Merck Patent GmbH, Germany
 SOURCE: Ger. Offen., 78 pp.
 CODEN: GWXXBX
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10315569	A1	20041014	DE 2003-10315569	20030405
AU 2004228124	A1	20041021	AU 2004-228124	20040310
CA 2521227	A1	20041021	CA 2004-2521227	20040310
WO 2004089932	A1	20041021	WO 2004-EP2453	20040310
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1611122	A1	20060104	EP 2004-718926	20040310
EP 1611122	B1	20070613		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK				
BR 2004008986	A	20060328	BR 2004-8986	20040310
CN 1768052	A	20060503	CN 2004-80008603	20040310
JP 2006522039	T	20060928	JP 2006-504620	20040310
AT 364601	T	20070715	AT 2004-718926	20040310
ES 2287710	T3	20071216	ES 2004-718926	20040310
US 20070010531	A1	20070111	US 2005-552064	20051005
PRIORITY APPLN. INFO.:			DE 2003-10315569	A 20030405
			WO 2004-EP2453	W 20040310
OTHER SOURCE(S):		MARPAT 141:332186		
GI				



AB Title compds. [I; R₁ = H, A, halo, (CH₂)_nAr, cycloalkyl, CF₃, NO₂, cyano, C(NH)NOH, OCF₃; R₂ = (CH₂)_nHet, (CH₂)_nAr, cycloalkyl, CF₃; R₃, R₄ = H, (CH₂)_nCO₂R₅, (CH₂)_nCOHet, CHO, (CH₂)_nOR₅, (CH₂)_nHet, CH:NOA, etc.; R₅ = H, A; A = alkyl, alkoxy, alkenyl, alkoxyalkyl; Ar = (substituted) Ph; Het = (aromatic) mono- or bicyclic heterocyclyl, heteroatom-containing organic residue; X

= N, CH; with provisos], were prepared Thus, [1-(4'-fluorobiphen-4-yl)-5-furan-2-yl-1H-pyrazol-4-ylmethyl]methyl(1-methylpyrrolidin-3-yl)amine showed 5-HT_{2A} activity with IC₅₀ = 5.14E-10.

=>

Connecting via Winsock to STN

Welcome to STN International! Enter x:X

LOGINID:ssptacrs1614

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page for STN Seminar Schedule - N. America
NEWS	2	DEC 01	ChemPort single article sales feature unavailable
NEWS	3	FEB 02	Simultaneous left and right truncation (SLART) added for CERAB, COMPUAB, ELCOM, and SOLIDSTATE
NEWS	4	FEB 02	GENBANK enhanced with SET PLURALS and SET SPELLING
NEWS	5	FEB 06	Patent sequence location (PSL) data added to USGENE
NEWS	6	FEB 10	COMPENDEX reloaded and enhanced
NEWS	7	FEB 11	WTEXTILES reloaded and enhanced
NEWS	8	FEB 19	New patent-examiner citations in 300,000 CA/CAPLUS patent records provide insights into related prior art
NEWS	9	FEB 19	Increase the precision of your patent queries -- use terms from the IPC Thesaurus, Version 2009.01
NEWS	10	FEB 23	Several formats for image display and print options discontinued in USPATFULL and USPAT2
NEWS	11	FEB 23	MEDLINE now offers more precise author group fields and 2009 MeSH terms
NEWS	12	FEB 23	TOXCENTER updates mirror those of MEDLINE - more precise author group fields and 2009 MeSH terms
NEWS	13	FEB 23	Three million new patent records blast AEROSPACE into STN patent clusters
NEWS	14	FEB 25	USGENE enhanced with patent family and legal status display data from INPADOCDB
NEWS	15	MAR 06	INPADOCDB and INPAFAMDB enhanced with new display formats
NEWS	16	MAR 11	EPFULL backfile enhanced with additional full-text applications and grants
NEWS	17	MAR 11	ESBIOBASE reloaded and enhanced
NEWS	18	MAR 20	CAS databases on STN enhanced with new super role for nanomaterial substances
NEWS	19	MAR 23	CA/CAPLUS enhanced with more than 250,000 patent equivalents from China
NEWS	20	MAR 30	IMSPATENTS reloaded and enhanced
NEWS	21	APR 03	CAS coverage of exemplified prophetic substances enhanced
NEWS	22	APR 07	STN is raising the limits on saved answers
NEWS	23	APR 24	CA/CAPLUS now has more comprehensive patent assignee information
NEWS	24	APR 26	USPATFULL and USPAT2 enhanced with patent assignment/reassignment information
NEWS	25	APR 28	CAS patent authority coverage expanded

NEWS 26 APR 28 ENCOMPLIT/ENCOMPLIT2 search fields enhanced
 NEWS 27 APR 28 Limits doubled for structure searching in CAS
 REGISTRY
 NEWS 28 MAY 08 STN Express, Version 8.4, now available
 NEWS 29 MAY 11 STN on the Web enhanced
 NEWS 30 MAY 11 BEILSTEIN substance information now available on
 STN Easy
 NEWS 31 MAY 14 DGENE, PCTGEN and USGENE enhanced with increased
 limits for exact sequence match searches and
 introduction of free HIT display format
 NEWS 32 MAY 15 INPADOCDB and INPAFAMDB enhanced with Chinese legal
 status data

NEWS EXPRESS MAY 26 09 CURRENT WINDOWS VERSION IS V8.4,
 AND CURRENT DISCOVER FILE IS DATED 06 APRIL 2009.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
 NEWS LOGIN Welcome Banner and News Items

Enter NEWS followed by the item number or name to see news on that
 specific topic.

All use of STN is subject to the provisions of the STN customer
 agreement. This agreement limits use to scientific research. Use
 for software development or design, implementation of commercial
 gateways, or use of CAS and STN data in the building of commercial
 products is prohibited and may result in loss of user privileges
 and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 11:03:14 ON 27 MAY 2009

=> file registry

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.22	0.22

FILE 'REGISTRY' ENTERED AT 11:03:27 ON 27 MAY 2009
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2009 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
 provided by InfoChem.

STRUCTURE FILE UPDATES: 25 MAY 2009 HIGHEST RN 1149058-00-3
 DICTIONARY FILE UPDATES: 25 MAY 2009 HIGHEST RN 1149058-00-3

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 9, 2009.


Please note that search-term pricing does apply when
 conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and
 predicted properties as well as tags indicating availability of
 experimental property data in the original document. For information
 on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stndoc/properties.html>

=>

Uploading C:\Program Files\STNEXP\Queries\10552065a.str



chain nodes :

12

ring nodes :

1 2 3 4 5 6 7 8 9 10 11

chain bonds :

5-7 11-12

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-11 8-9 9-10 10-11

exact/norm bonds :

5-7 7-8 7-11 8-9 9-10 10-11

exact bonds :

11-12

normalized bonds :

1-2 1-6 2-3 3-4 4-5 5-6

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom

11:Atom 12:CLASS

L1 STRUCTURE UPLOADED

=> s l1 sss ful

FULL SEARCH INITIATED 11:03:43 FILE 'REGISTRY'

FULL SCREEN SEARCH COMPLETED - 828790 TO ITERATE

100.0% PROCESSED 828790 ITERATIONS


434431 ANSWERS

SEARCH TIME: 00.00.12

L2 434431 SEA SSS FUL L1

=>

Uploading C:\Program Files\STNEXP\Queries\10552065b.str



```

ring nodes :
1  2  3  4  5  6  7  8  9  10  11  12  13  14  15  16  17
chain bonds :
5-7  11-15
ring bonds :
1-2  1-6  2-3  3-4  4-5  5-6  7-8  7-11  8-9  9-10  10-11  12-13  12-17  13-14
14-15  15-16  16-17
exact/norm bonds :
5-7  7-8  7-11  8-9  9-10  10-11
exact bonds :
11-15
normalized bonds :
1-2  1-6  2-3  3-4  4-5  5-6  12-13  12-17  13-14  14-15  15-16  16-17

```

```

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:Atom

```

L3 STRUCTURE UPLOADED

```

=> s l3 fam ful
FULL SEARCH INITIATED 11:05:32 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED -       3064 TO ITERATE

```

```

100.0% PROCESSED       3064 ITERATIONS                   2 ANSWERS
SEARCH TIME: 00.00.01

```

L4 2 SEA FAM FUL L3

```

=> file caplus
COST IN U.S. DOLLARS                   SINCE FILE       TOTAL
                                      ENTRY       SESSION
FULL ESTIMATED COST                   260.17       260.39

```

```

FILE 'CAPLUS' ENTERED AT 11:05:41 ON 27 MAY 2009
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

```

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

```

FILE COVERS 1907 - 27 May 2009  VOL 150 ISS 22
FILE LAST UPDATED: 26 May 2009  (20090526/ED)
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Feb 2009
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Feb 2009

```

Caplus now includes complete International Patent Classification (IPC) reclassification data for the third quarter of 2008.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate

=> s 14

L5 38 L4

=> d 15 ibib abs 1-38

L5 ANSWER 1 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2009:338987 CAPLUS

DOCUMENT NUMBER: 150:329777

TITLE: Copper-catalyzed C-H bond arylation of heterocyclic compounds and electron deficient arenes with aryl halides

INVENTOR(S): Daugulis, Olafs; Do, Hien-Quang

PATENT ASSIGNEE(S): The University of Houston System, USA

SOURCE: U.S. Pat. Appl. Publ., 112pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20090076266	A1	20090319	US 2008-208286	20080910
PRIORITY APPLN. INFO.:			US 2007-971466P	P 20070911

OTHER SOURCE(S): CASREACT 150:329777

AB The present invention is a one-step method for efficiently converting carbon-hydrogen bonds into carbon-carbon bonds using a combination of aryl halides, a substrate, and a copper salt as catalyst. Thus, e.g., 2-phenylbenzoxazole was prepared in 93% yield by reacting benzoxazole with iodobenzene in the presence of a catalytic amount of copper(I) iodide and base in DMF with heating. This method allows faster introduction of complex mol. entities, a process that would otherwise require many more steps. This invention is particularly relevant for the organic synthesis of complex mols. such as, but not limited to, pharmacophores and explosives.

L5 ANSWER 2 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:1383562 CAPLUS

DOCUMENT NUMBER: 149:555078

TITLE: The Stille reaction

AUTHOR(S): Farina, Vittorio; Krishnamurthy, Venkat; Scott, William J.

CORPORATE SOURCE: Boehringer Ingelheim Pharmaceuticals, Ridgefield, CT, USA

SOURCE: Organic Reactions (Hoboken, NJ, United States) (1997), 50, No pp. given

CODEN: ORHNBA

URL: <http://www3.interscience.wiley.com/cgi-bin/mrwhome/107610747/HOME>

PUBLISHER: John Wiley & Sons, Inc.

DOCUMENT TYPE: Journal; General Review; (online computer file)

LANGUAGE: English

OTHER SOURCE(S): CASREACT 149:555078

AB A review of the article The Stille reaction.

L5 ANSWER 3 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2008:1243395 CAPLUS

DOCUMENT NUMBER: 149:534179
TITLE: A General Method for Copper-Catalyzed Arylation of Arene C-H Bonds
AUTHOR(S): Do, Hien-Quang; Khan, Rana M. Kashif; Daugulis, Olafs
CORPORATE SOURCE: Department of Chemistry, University of Houston, Houston, TX, 77204-5003, USA
SOURCE: Journal of the American Chemical Society (2008), 130(45), 15185-15192
CODEN: JACSAT; ISSN: 0002-7863
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
AB A general method for copper-catalyzed arylation of sp² C-H bonds with pK_a's below 35 has been developed. The method employs aryl halide as the coupling partner, lithium alkoxide or K₃PO₄ base, and DMF, DMPU, or mixed DMF/xylenes solvent. A variety of electron-rich and electron-poor heterocycles such as azoles, caffeine, thiophenes, benzofuran, pyridine oxides, pyridazine, and pyrimidine can be arylated. Furthermore, electron-poor arenes possessing at least two electron-withdrawing groups on a benzene ring can also be arylated. Two arylcopper-phenanthroline complex intermediates were independently synthesized.
REFERENCE COUNT: 77 THERE ARE 77 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 4 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2007:554503 CAPLUS
DOCUMENT NUMBER: 147:143327
TITLE: Regioselective microwave-assisted synthesis of substituted pyrazoles from ethynyl ketones
AUTHOR(S): Bagley, Mark C.; Lubinu, M. Caterina; Mason, Christopher
CORPORATE SOURCE: School of Chemistry, Cardiff University, Cardiff, CF10 3AT, UK
SOURCE: Synlett (2007), (5), 704-708
CODEN: SYNLES; ISSN: 0936-5214
PUBLISHER: Georg Thieme Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 147:143327
AB Reaction of α,β -ethynyl ketones and hydrazine derivs. gives 1,3- and 1,5-disubstituted pyrazoles in good yield. Microwave irradiation in concentrated HCl/MeOH (1.5% volume/volume), with concurrent cooling at sub-ambient temps. or at 120°, for 30 or 2 min, resp., facilitates rapid heterocyclization and preferentially gives the 1,3-disubstituted regioisomer.
REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 5 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2007:220868 CAPLUS
DOCUMENT NUMBER: 146:304597
TITLE: Electroluminescent organometallic complexes, organic electroluminescent devices, and displays and lightings using them
INVENTOR(S): Oshiyama, Tomohiro; Yasukawa, Noriko; Kato, Eisaku
PATENT ASSIGNEE(S): Konica Minolta Holdings, Inc., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 52pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2007051243	A	20070301	JP 2005-238536	20050819
PRIORITY APPLN. INFO.:			JP 2005-238536	20050819
OTHER SOURCE(S):	MARPAT	146:304597		
GI				

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB The electroluminescent organometallic complexes have partial structure of I (R01-R07 = H, substituent; ≥ 1 of R01, R03-R05, R07 = aromatic heterocyclic group; M01 = Group 8-10 metal). Also claimed are organometallic complexes having partial structures of I, wherein R01 or R05 represents an aromatic heterocyclic group. Also claimed are organometallic complexes of II (R2-R27 = H, substituent; R24 = aromatic heterocyclic group; X1-L1-X2 = bidentate ligand; X1, X2 = C, N; L1 = group of atoms; n2 = 1-3; m2 = 0-2; n2 + m2 = 2, 3; M21 = Group 8-10 metal). Organic electroluminescent (EL) devices with high emission efficiency and long life are provided with this invention.

L5 ANSWER 6 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:529471 CAPLUS

DOCUMENT NUMBER: 145:188781

TITLE: Palladium-catalyzed C-N bond formation: synthesis of 1-aryl-1H-pyrazoles from β -bromovinyl aldehydes and arylhydrazines

AUTHOR(S): Cho, Chan Sik; Patel, Daksha B.

CORPORATE SOURCE: Research Institute of Industrial Technology, Kyungpook National University, Taegu, 702-701, S. Korea

SOURCE: Tetrahedron (2006), 62(26), 6388-6391

CODEN: TETRAB; ISSN: 0040-4020

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 145:188781

AB 1-Aryl-1H-pyrazoles and fused arylpyrazoles are prepared in 20-79% yields by cyclocondensation of arylhydrazines with cyclic and acyclic β -bromo- α,β -unsatd. aldehydes in the presence of palladium acetate, a diphosphine such as 1,1'-bis(diphenylphosphino)ferrocene (dppf), and sodium tert-butoxide in toluene at 125°.

REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 7 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2006:202383 CAPLUS

DOCUMENT NUMBER: 145:489165

TITLE: Iodine(III) mediated synthesis of new 5-aryl-3-(4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl)-1-phenylpyrazoles from dehydrogenation of 5-aryl-3-(4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl)-1-phenylpyrazolines

AUTHOR(S): Prakash, Om; Kumar, Ajay; Kinger, Mayank; Singh, Shiv P.

CORPORATE SOURCE: Department of Chemistry, Kurukshetra University, Kurukshetra, 136 119, India

SOURCE: Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry (2006),

45B(2), 456-460
CODEN: IJSBDB; ISSN: 0376-4699
PUBLISHER: National Institute of Science Communication and
Information Resources
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 145:489165
AB 3-Cinnamoyl-4-hydroxy-6-methyl-2-pyrones (chalcone analogs of DHA) on
condensation with PhN₂H₃ in EtOH, yield
5-aryl-3-(4-hydroxy-6-methyl-2-oxo-2H-pyran-3-yl)-1-phenylpyrazolines
which undergo smooth dehydrogenation to the corresponding pyrazoles in
good yield upon treatment with iodobenzene diacetate (IBD).
REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 8 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2005:1118021 CAPLUS
DOCUMENT NUMBER: 144:51499
TITLE: Regioselective Synthesis of
1-Aryl-3,4-substituted/annulated-5-
(methylthio)pyrazoles and
1-Aryl-3-(methylthio)-4,5-substituted/annulated
Pyrazoles
AUTHOR(S): Peruncheralathan, S.; Khan, T. A.; Ila, H.; Junjappa,
H.
CORPORATE SOURCE: Department of Chemistry, Indian Institute of
Technology, Kanpur, 208016, India
SOURCE: Journal of Organic Chemistry (2005), 70(24),
10030-10035
CODEN: JOCEAH; ISSN: 0022-3263
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 144:51499
AB Highly efficient and regioselective synthesis of
1-aryl-3,4-substituted/annulated-5-(methylthio)pyrazoles and
1-aryl-3-(methylthio)-4,5-substituted/annulated pyrazoles has been
reported via cyclocondensation of arylhydrazines with either
 α -oxoketene dithioacetals or β -oxo dithio esters.
REFERENCE COUNT: 77 THERE ARE 77 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 9 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2002:855864 CAPLUS
DOCUMENT NUMBER: 139:214344
TITLE: Product class 1: pyrazoles
AUTHOR(S): Stanovnik, B.; Svete, J.
CORPORATE SOURCE: Faculty of Chemistry and Chemical Technology, Division
of Organic Chemistry, Ljubljana, 61000, Slovenia
SOURCE: Science of Synthesis (2002), 12, 15-225
CODEN: SSCYJ9
PUBLISHER: Georg Thieme Verlag
DOCUMENT TYPE: Journal; General Review
LANGUAGE: English
AB A review. Methods for preparing pyrazoles are reviewed including
cyclization, ring transformation, aromatization and substituent
modifications.
REFERENCE COUNT: 909 THERE ARE 909 CITED REFERENCES AVAILABLE FOR
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

L5 ANSWER 10 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 2001:321317 CAPLUS
DOCUMENT NUMBER: 135:122152
TITLE: A complete model for the prediction of ¹H- and ¹³C-NMR chemical shifts and torsional angles in phenyl-substituted pyrazoles
AUTHOR(S): Carrillo, J. R.; Cossio, F. P.; Diaz-Ortiz, A.; Gomez-Escalonilla, M. J.; de la Hoz, A.; Lecea, B.; Moreno, A.; Prieto, P.
CORPORATE SOURCE: Facultad de Quimica, Universidad de Castilla-La Mancha, Ciudad Real, 13071, Spain
SOURCE: Tetrahedron (2001), 57(19), 4179-4187
CODEN: TETRAB; ISSN: 0040-4020
PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
AB ¹H- and ¹³C-NMR spectra of a number of N-phenyl- and C-phenylpyrazole derivs. have been obtained. The parameter most susceptible to changes in the dihedral angle is the difference $\delta_{\text{meta-C}} - \delta_{\text{ortho-C}}$. Values for this parameter have been determined and its usefulness for conformational studies of phenyl-substituted pyrazoles has been demonstrated. A correlation between torsional angles calculated by mol. mechanics and differences in ¹³C chemical shifts of the ortho and meta carbon atoms of the Ph groups in 29 N-phenyl-substituted pyrazole derivs. and 11 C-phenyl-substituted pyrazole derivs. has been found. For the N-phenyl-substituted derivs. a correlation between torsional angles and $\delta_{\text{meta-H}} - \delta_{\text{ortho-H}}$ values has also been demonstrated. In all cases good correlations between angles and differences in chemical shifts were observed
REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 11 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 2000:500201 CAPLUS
DOCUMENT NUMBER: 133:281725
TITLE: A New Germanium-Based Linker for Solid Phase Synthesis of Aromatics: Synthesis of a Pyrazole Library
AUTHOR(S): Spivey, Alan C.; Diaper, Christopher M.; Adams, Harry; Rudge, Andrew J.
CORPORATE SOURCE: Department of Chemistry, University of Sheffield, Yorkshire, S3 7HF, UK
SOURCE: Journal of Organic Chemistry (2000), 65(17), 5253-5263
CODEN: JOCEAH; ISSN: 0022-3263
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 133:281725
AB An efficient synthesis of chlorogermane linker 4-HOC₆H₄CH₂CH₂GeMe₂Cl (I) is described. Economic introduction of germanium into this linker is accomplished by insertion of dichlorogermylene [from germanium(IV) chloride] into the homobenzylic C-Cl bond of 4-HOC₆H₄CH₂CH₂Cl. Using linker I, transmetalation with lithiated 4-acetophenone, 3-acetophenone, and 4-methoxybiphenyl followed by Mitsunobu-type coupling to Argogel gives functionalized resins. Treatment of resin-bound 4-HOC₆H₄CH₂CH₂GeMe₂Cl with TFA, ICl, Br₂, or NCS effects clean ipso-degermylation releasing 4-MeOC₆H₄CH₂CH₂GeMe₂Cl [R = H, I, Br, Cl]. Resin-bound 4-HOC₆H₄CH₂CH₂GeMe₂Cl [R₁ = 4-Ac, 3-Ac] are employed for the parallel synthesis of a library of pyrazoles by enaminone formation (using Bredereck's reagent), condensative ring-closure (using a series of monosubstituted hydrazines), and cleavage (using TFA and Br₂). Anal. of this library reveals the influence of the hydrazine substituent on both the regioselectivity of ring-closure and the propensity for electrophilic substitution at the 4-position of the pyrazoles during ipso-degermylation

cleavage.

REFERENCE COUNT: 96 THERE ARE 96 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 12 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1997:57955 CAPLUS
DOCUMENT NUMBER: 126:211777
ORIGINAL REFERENCE NO.: 126:40951a
TITLE: Substituent effects on the ¹⁵N NMR parameters of azoles
AUTHOR(S): Claramunt, Rosa Maria; Sanz, Dionisia; Lopez, Concepcion; Jimenez, Jose Antonio; Jimeno, Maria Luisa; Elguero, Jose; Fruchier, Alain
CORPORATE SOURCE: Departamento de Quimica Organica y Biologia, Facultad de Ciencias, UNED, Madrid, E-28040, Spain
SOURCE: Magnetic Resonance in Chemistry (1997), 35(1), 35-75
CODEN: MRCHEG; ISSN: 0749-1581
PUBLISHER: Wiley
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The ¹⁵N chemical shifts and a large collection of coupling consts. pertaining to azoles have been gathered from the literature. To complete this collection and to check some anomalies, the spectra of 14 compds. in several solvents were recorded again and 31 compds. were studied for the first time; in all, data for 420 compds. (pyrroles, imidazoles, pyrazoles, triazoles, tetrazoles, indoles, benzimidazoles, indazoles, benzotriazoles and carbazoles) are reported. Additive models are used to discuss the substituent chemical shifts.

REFERENCE COUNT: 52 THERE ARE 52 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 13 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN


ACCESSION NUMBER: 1992:425806 CAPLUS
DOCUMENT NUMBER: 117:25806
ORIGINAL REFERENCE NO.: 117:4639a,4642a
TITLE: Carbon-13 chemical shifts and proton-13C coupling constants of N-phenyl-, N-p-fluorophenyl- and N-o-nitrophenylpyrazoles
AUTHOR(S): Begtrup, Mikael; Vedsoe, Per; Cabildo, Pilar; Claramunt, Rosa Maria; Elguero, Jose; Meutermans, Wim
CORPORATE SOURCE: Dep. Org. Chem., R. Dan. Sch. Pharm., Copenhagen, DK-2100, Den.
SOURCE: Magnetic Resonance in Chemistry (1992), 30(5), 455-9
CODEN: MRCHEG; ISSN: 0749-1581
DOCUMENT TYPE: Journal
LANGUAGE: English

AB The ¹³C chemical shifts and some ¹H-¹³C coupling consts. of twelve N-arylpyrazoles are reported. The assignments were made by using the effects of a fluorine substituent and two-dimensional techniques.

L5 ANSWER 14 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1988:528890 CAPLUS
DOCUMENT NUMBER: 109:128890
ORIGINAL REFERENCE NO.: 109:21473a,21476a
TITLE: Reaction of 2-dimethylaminomethylene-1,3-diones with dinucleophiles. VI. Synthesis of ethyl or methyl 1,5-disubstituted 1H-pyrazole-4-carboxylates
AUTHOR(S): Menozzi, Giulia; Mosti, Luisa; Schenone, Pietro
CORPORATE SOURCE: Ist. Sci. Farm., Univ. Genoa, Genoa, 16132, Italy
SOURCE: Journal of Heterocyclic Chemistry (1987), 24(6), 1669-75
CODEN: JHTCAD; ISSN: 0022-152X


DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 109:128890
 GI



AB Reaction of $\text{RCOCH}_2\text{CO}_2\text{R}^1$ ($\text{R} = \text{alkyl, Ph, PhCH}_2$; $\text{R}^1 = \text{Me, Et}$) with N,N-dimethylformamide di-Me acetal gave, generally in excellent yields, $\text{RCOC}(:\text{CHNMe}_2)\text{CO}_2\text{R}^1$ (I) which reacted with phenylhydrazine to afford the esters (II) of 5-substituted 1-phenyl-1H-pyrazole-4-carboxylic acids in high yields. Ester II were hydrolyzed to 5-substituted 1-phenyl-1H-pyrazole-4-carboxylic acids which were converted by heating to 5-substituted 1-phenyl-1H-pyrazoles III in excellent yields. Reaction of I with methylhydrazine afforded in general a mixture of 3- and 5-substituted Et 1-methyl-1H-pyrazole-4-carboxylates with the exception of I ($\text{R} = \text{PhCH}_2$, $\text{R}^1 = \text{Me}$), which gave in high yield Me 5-benzyl-1-methyl-1H-pyrazole-4-carboxylate, which was hydrolyzed to the corresponding pyrazolecarboxylic acid. This afforded by heating 5-benzyl-1-methyl-1H-pyrazole in quant. yield.

L5 ANSWER 15 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN


ACCESSION NUMBER: 1988:186642 CAPLUS
 DOCUMENT NUMBER: 108:186642
 ORIGINAL REFERENCE NO.: 108:30667a,30670a
 TITLE: Oxidative cyclization of arylhydrazones of chalcones and benzalacetones to pyrazoles by thianthrene cation radical
 AUTHOR(S): Kovelesky, Albert C.; Shine, Henry J.
 CORPORATE SOURCE: Dep. Chem. Biochem., Texas Tech Univ., Lubbock, TX, 79409, USA
 SOURCE: Journal of Organic Chemistry (1988), 53(9), 1973-9
 CODEN: JOCEAH; ISSN: 0022-3263
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 108:186642
 GI



AB Phenyl-, (p-nitrophenyl)-, and (2,4-dinitrophenyl)hydrazones of chalcone (benzalacetophenone), benzalacetone, and of some of their derivs. undergo oxidative cyclization in reaction with thianthrene cation radical perchlorate. The products are, e.g., 1,3,5-triaryl- (I; $\text{R} = \text{Ph}$; $\text{R}^1, \text{R}^2 = \text{H, NO}_2$) and 3-methyl-1,5-diarylpyrazoles (I; $\text{R} = \text{Me}$; $\text{R}^1, \text{R}^2 = \text{H, NO}_2$) and



are formed in excellent yields. Cyclization appears to occur by way of the arylhydrazone cation radical and not via the preliminary, acid-catalyzed formation of the corresponding pyrazoline.

L5 ANSWER 16 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 1985:596029 CAPLUS
DOCUMENT NUMBER: 103:196029
ORIGINAL REFERENCE NO.: 103:31589a,31592a
TITLE: The reactions of isothiazolium salts with nitrogen nucleophilic reagents
AUTHOR(S): Hassan, Mohamed E.; Magraby, M. A.; Aziz, Magda A.
CORPORATE SOURCE: Chem. Dep., Aswan Univ., Aswan, Egypt
SOURCE: Tetrahedron (1985), 41(10), 1885-91
CODEN: TETRAB; ISSN: 0040-4020
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 103:196029
GI




AB Isothiazolium salts I (R = R1 = H, R2 = Ph, R3 = Me, Ph, X = ClO4; R = H, R1 = SMe, R2 = Ph, R3 = Me, X = ClO4; R = H, R1 = Ph, R2 = SMe, R3 = Me, X = iodide; R = R2 = H, R1 = Ph, R3 = Me; X = ClO4; R = Ph, C6H4Me-p, R1 = R2 = H, R3 = Me, X = ClO4) reacted with a number of N nucleophiles including NH3, PhNHNH2, H2NOH and PhCH2NH2. The products obtained suggest that the position of initial nucleophilic attack is at the S of the heterocyclic cation.

L5 ANSWER 17 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
ACCESSION NUMBER: 1984:530627 CAPLUS
DOCUMENT NUMBER: 101:130627
ORIGINAL REFERENCE NO.: 101:19873a,19876a
TITLE: New synthesis of pyrazole and isoxazole derivatives
AUTHOR(S): Molina, P.; Fresneda, P. M.
CORPORATE SOURCE: Fac. Cienc., Univ. Murcia, Murcia, Spain
SOURCE: Journal of Heterocyclic Chemistry (1984), 21(2), 461-4
CODEN: JHTCAD; ISSN: 0022-152X
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 101:130627
GI



AB 1-Phenylpyrazoles I (R = H, Me, NO₂, MeO, Br, Cl) and isoxazoles II were prepared by treating ketimines III (R₁ = Me) with (MeO)₂CHNMe₂ to give enaminimines III (R₁ = CH:CHNMe₂) which were cyclocondensed with PhNHNH₂ or H₂NOH.HCl.


L5 ANSWER 18 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
 ACCESSION NUMBER: 1980:604527 CAPLUS
 DOCUMENT NUMBER: 93:204527
 ORIGINAL REFERENCE NO.: 93:32633a,32636a
 TITLE: Reaction of activated methyl groups with
 N,N-dimethylformamide dialkyl acetals
 AUTHOR(S): Tisler, M.; Stanovnik, B.; Vercek, B.
 CORPORATE SOURCE: Dep. Chem., Univ. Ljubljana, Ljubljana, 61000,
 Yugoslavia
 SOURCE: Vestnik Slovenskega Kemijskega Drustva (1980), 27(1),
 65-72
 CODEN: VSKDAA; ISSN: 0560-3110
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 93:204527
 GI



AB RCOCH:CHNMe₂ (I, R = 2-pyridyl, 3-pyridyl, 2-O₂NC₆H₄) were prepared in 49-73% yield by treating RAc with Me₂NCH(OMe)₂. The pyrazoles II (R₁ = H, Ph) were obtained by treating I with R₁NHNH₂. II (R = R₁ = Ph) was similarly obtained from BzCH:CHNMe₂. The quinazoline oxide III was prepared in 56% yield by treating 2-H₂NC₆H₄CMe:NOH with Me₂NCH(OEt)₂.

L5 ANSWER 19 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
 ACCESSION NUMBER: 1980:41480 CAPLUS
 DOCUMENT NUMBER: 92:41480
 ORIGINAL REFERENCE NO.: 92:6913a,6916a
 TITLE: Reactivity of α-halogenated imino compounds.
 Part XVIII. Reactivity of

N-aryl- α,α -dichlorinated arylketimines
 AUTHOR(S): De Kimpe, Norbert; Verhe, Roland; De Buyck, Laurent;
 CORPORATE SOURCE: Tukiman, Sunari; Schamp, Niceas
 SOURCE: Lab. Org. Chem., State Univ. Gent, Ghent, Belg.
 Tetrahedron (1979), 35(6), 789-98
 CODEN: TETRAB; ISSN: 0040-4020
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 92:41480
 AB E-RCCl₂C(C₆H₄R₁-4):NC₆H₄R₂ (I; R = alkyl, R₁ = H, Br, R₂ = H, Me, OMe) were prepared by chlorination of E-RCH₂C(C₆H₄R₁-4):NC₆H₄R₂ by N-chlorosuccinimide in CCl₄. I (R = Me) (R₁ = H, R₂ = H, p-Me, m-Me, p-OMe; R₁ = Br, R₂ = H) with NaOMe/MeOH, followed by acid hydrolysis, gave 4-R₁C₆H₄COCOMe, 4-R₁C₆H₄COC(OMe):CH₂, and 4-R₁C₆H₄COCH:CHNHC₆H₄R₂, formation of the latter formally involving a migration of N from C-1 to C-3. More highly substituted I with NaOMe/MeOH gave mainly α -chloro- α,β -unsatd. ketones. With long-chain I, a formal γ -functionalization was observed. Reaction mechanisms are discussed in detail.
 L5 ANSWER 20 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN
 ACCESSION NUMBER: 1978:190700 CAPLUS
 DOCUMENT NUMBER: 88:190700
 ORIGINAL REFERENCE NO.: 88:29989a
 TITLE: 2-(Acylmethyl)-1,3,4-dioxazoles by ketovinylation of hydroxamic acids
 AUTHOR(S): Schroth, Werner; Peters, Olaf
 CORPORATE SOURCE: Sek. Chem., Martin-Luther-Univ., Halle/Saale, Ger. Dem. Rep.
 SOURCE: Zeitschrift fuer Chemie (1978), 18(2), 57-8
 CODEN: ZECEAL; ISSN: 0044-2402
 DOCUMENT TYPE: Journal
 LANGUAGE: German
 OTHER SOURCE(S): CASREACT 88:190700
 GI



AB Hydroxamic acids RCONHOH reacted with β -chlorovinyl ketones R₁COCH:CHCl to give 21-96% of the dioxazoles I (R = Ph, styryl, 4-O₂NC₆H₄, CH₂Cl; R₁ = Me, Me₂CH, Ph, 4-ClC₆H₄, 4-O₂NC₆H₄, o-tolyl). I were hydrolyzed by alc. alkali to hydroxamic acids and the corresponding β -dicarbonyl compds. Reaction of I with 2,4-R₂(O₂N)C₆H₃NHNH₂ (R₂ = H, NO₂) gave the pyrazoles II (R₁ and R₂ as above).
 L5 ANSWER 21 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1976:17219 CAPLUS
 DOCUMENT NUMBER: 84:17219
 ORIGINAL REFERENCE NO.: 84:2851a, 2854a
 TITLE: Intermediates isolated during the synthesis of 1,3-

and 1,5-diphenylpyrazoles
AUTHOR(S): Rull, Thomas; Le Strat, Georges
CORPORATE SOURCE: Cent. Rech. ATO Chim., Orsay, Fr.
SOURCE: Bulletin de la Societe Chimique de France (1975),
(5-6, Pt. 2), 1375-9
CODEN: BSCFAS; ISSN: 0037-8968

DOCUMENT TYPE: Journal
LANGUAGE: French
OTHER SOURCE(S): CASREACT 84:17219

GI For diagram(s), see printed CA Issue.

AB BzCH:CHNHNHPh, and pyrazolines I (R = Ph, R1 = H; R = H, R1 = Ph) were isolated as intermediates in the reaction of BzCH2CHO with PhNHNH2 to give title diphenylpyrazoles. Reaction of BzCH:CHOBz with PhNHNH2 gave BzCH:CHNPhNH2 as the intermediate to 1,3-diphenylpyrazole. Reaction of BzCH2CHO with PhNHNHCHO gave BzCH:CHNPhNHCHO, which gave 1,3- and 1,5-diphenylpyrazoles in 77.5:22.5 ratio and 82% yield.

L5 ANSWER 22 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1971:420283 CAPLUS

DOCUMENT NUMBER: 75:20283

ORIGINAL REFERENCE NO.: 75:3247a,3250a

TITLE: Sulfurated organic compounds. VIII. Reaction of phenylhydrazine with 3-aryl-1,2-dithiolylum bisulfates and 3,5-diaryl-1,2-dithiolylum perchlorates. 3-Aryl-1-phenylpyrazoles, 5-aryl-1-phenylpyrazoles, and 3,5-diaryl-1-phenylpyrazoles

AUTHOR(S): Bergeon, Marie T.; Metayer, Claire; Quiniou, Herve

CORPORATE SOURCE: Fac. Sci., Nantes, Fr.

SOURCE: Bulletin de la Societe Chimique de France (1971), (3), 917-24
CODEN: BSCFAS; ISSN: 0037-8968

DOCUMENT TYPE: Journal

LANGUAGE: French

GI For diagram(s), see printed CA Issue.

AB PhNHNH2 (I) and 3-aryl dithioles (II) reacted to give 2 pyrazole isomers (III) which were separated by Al2O3 chromatog. Thus, II (R = Ph, R1 = H, X = HSO4) reacted with I in EtOH to give III (R = Ph, R1 = H) (major) and III (R = H, R1 = Ph). II (R = R1 = Ph, X = ClO4) reacted with I to give 60% III (R = R1 = Ph). II (R = Ph, R1 = p-ClC6H4, X = ClO4) when treated similarly gave the isomeric III (R = Ph, R1 = p-ClC6H4) (major) and III (R = p-ClC6H4, R1 = Ph). The structure of pyrazoles was determined from chemical anal. and NMR spectra.

L5 ANSWER 23 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1971:99117 CAPLUS

DOCUMENT NUMBER: 74:99117

ORIGINAL REFERENCE NO.: 74:16133a,16136a

TITLE: Investigations on pyrazole derivatives. V. Uv absorption spectra of derivatives of 6H,7H-pyrazolo [3,2-b][1,2,4]thiazole and some pyrazole derivatives

AUTHOR(S): Dymek, Wojciech; Ryznerski, Zygmunt

CORPORATE SOURCE: Dep. Pharm. Chem., Med. Acad., Cracow, Pol.

SOURCE: Dissertationes Pharmaceuticae et Pharmacologicae (1970), 22(6), 419-25
CODEN: DPHFAK; ISSN: 0012-3870

DOCUMENT TYPE: Journal

LANGUAGE: English

GI For diagram(s), see printed CA Issue.

AB The uv absorption spectra of substituted 6H, 7H-pyrazolo[3,2-b][1,2,4]thiadiazoles (I) exhibited a bathochromic shift and a change in extinction coefficient in comparison with the spectra of the

corresponding pyrazole derivs. The most pronounced changes in the maxima positions are produced by substituents at position 7 in I and at position 4 in pyrazole. Substituents at other positions caused little change in the maxima position but did influence the intensity.

L5 ANSWER 24 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1970:132597 CAPLUS

DOCUMENT NUMBER: 72:132597

ORIGINAL REFERENCE NO.: 72:23735a,23738a

TITLE: Azoles. LXIII. Reactions of mono- and 1,2-disubstituted hydrazines with α -acetylenic carbonyl compounds

AUTHOR(S): Coispeau, Gerard; Elguero, Jose; Jacquier, Robert

CORPORATE SOURCE: Lab. Syn. Etude Phys.-Chim. Heterocycles Azotes, Fac. Sci., Montpellier, Fr.

SOURCE: Bulletin de la Societe Chimique de France (1970), (2), 689-96

CODEN: BSCFAS; ISSN: 0037-8968

DOCUMENT TYPE: Journal

LANGUAGE: French

AB The reactions of MeNHNH₂, PhNHNH₂, and 2,4-dinitrophenyl-hydrazine with α -acetylenic carbonyls such as HC.tplbond.CCH(OEt)₂ (i.e., acetals were used when the aldehydes were unstable), HC.tplbond.CAc, HC.tplbond.CBz, and PhC.tplbond.CAc gave N-substituted-pyrazoles. The reactions of MeNHNHPh with the same carbonyls in the presence of HI gave the corresponding pyrazolium iodides. Contrary to analogous condensations with β -diketones, the orientation of the reaction permits the preparation of certain isomers which are difficult to prepare

L5 ANSWER 25 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1968:512873 CAPLUS

DOCUMENT NUMBER: 69:112873

ORIGINAL REFERENCE NO.: 69:21119a,21122a

TITLE: Polarographic study of derivatives of Δ^2 -pyrazoline

AUTHOR(S): Shimanskaya, N. P.; Buryakovskaya, E. G.; Bezuglyi, V. D.; Tsukerman, S. V.

CORPORATE SOURCE: Vses. Nauch.-Issled. Inst. Monokrist., Kharkov, USSR

SOURCE: Zhurnal Obshchei Khimii (1968), 38(8), 1676-9

CODEN: ZOKHA4; ISSN: 0044-460X

DOCUMENT TYPE: Journal

LANGUAGE: Russian

AB Polarographic data are presented, mainly graphically, for 14 examples of 2-pyrazolines with a 1-phenyl substituent and substituents in 3- and 5-positions selected from Ph, H, 2-furyl, 2-selenophene-yl, p-dimethylaminophenyl, 2-thienyl, p-anisyl, 2,4-dimethoxyphenyl, Me, and p-ClC₆H₄. Substituents in the 3-position affect the half-wave potentials materially and the declining potentials follow the rise in electron-acceptor capability of these groups in order: 2-furyl, 2-thienyl, 2-selenophene-yl. Substituents in the 5-position exert only their inductive effects. The half-wave potentials were readily correlated with the absorption spectral long-wavelength maximum of each compound with the wavelength of the band declining linearly with rising neg. value of the half-wave potential.

L5 ANSWER 26 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1968:451379 CAPLUS

DOCUMENT NUMBER: 69:51379

ORIGINAL REFERENCE NO.: 69:9579a,9582a

TITLE: Molecular orbital calculations of pyrazoles. I. Alkyl- and aryl-pyrazoles

AUTHOR(S): Finar, I. L.

CORPORATE SOURCE: Northern Polytech., London, UK
SOURCE: Journal of the Chemical Society [Section] B: Physical Organic (1968), (7), 725-32
CODEN: JCSPAC; ISSN: 0045-6470

DOCUMENT TYPE: Journal
LANGUAGE: English

AB The longest $\pi \rightarrow \pi^*$ wave absorption band in the uv spectra of 40 alkyl- and arylpyrazoles was calculated by the simple L.C.A.O.M.O. method. With a suitable choice of parameters, a good correlation was obtained between the calculated and observed frequencies. Angles of twist were calculated for some sterically hindered pyrazoles, and the stabilities of some tautomeric forms of 1-unsubstituted pyrazoles and the basicities of a number of pyrazoles were examined. Electrophilic and homolytic substitution in pyrazole, 1-methyl- and 1-phenylpyrazole, and their corresponding conjugate acids were discussed in terms of reactivity indices. 49 references.

L5 ANSWER 27 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1968:402346 CAPLUS
DOCUMENT NUMBER: 69:2346
ORIGINAL REFERENCE NO.: 69:439a,442a
TITLE: Mass spectra and structure of organic compounds.
XXIV. Mass spectra of phenyl derivatives of pyrazole.

AUTHOR(S): Krasnoshchek, A. P.; Khmel'nitskii, R. A.; Polyakova, A. A.; Grandberg, I. I.

CORPORATE SOURCE: Mosk. Gos. Univ. im. Lomonosova, Moscow, USSR
SOURCE: Zhurnal Organicheskoi Khimii (1968), 4(4), 689-95
CODEN: ZORKAE; ISSN: 0514-7492

DOCUMENT TYPE: Journal
LANGUAGE: Russian
OTHER SOURCE(S): CASREACT 69:2346

GI For diagram(s), see printed CA Issue.

AB Mass spectra of the following pyrazoles (I) were obtained [substituent(s), position(s) in I given]: Ph, 1; Ph, 3 (5); Ph, Ph, 1, 3; Ph, Ph, 1, 5; Ph, Ph, 1, 4; Ph, Ph, 3, 5; Ph, Ph, Ph, 1, 3, 5; Ph, Ph, Ph, 3, 4, 5. The stability towards electron impact of I decreased with the number of C-Ph groups and it increased with the number of N-Ph groups. Migrations of Ph groups were observed. A 9-fluorenyl ion (II) was formed from I by an intramol. condensation-elimination mechanism.

L5 ANSWER 28 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1967:516333 CAPLUS
DOCUMENT NUMBER: 67:116333
ORIGINAL REFERENCE NO.: 67:21890h,21891a
TITLE: 1,3-Dipolar cycloadditions. XXXIII. Differences in the reactivity of substituted nitrilimines
AUTHOR(S): Huisgen, Rolf; Adelsberger, Klaus; Aufderhaar, Ernst; Knupfer, Hans; Wallbillich, Guenter
CORPORATE SOURCE: Univ. Munich, Munich, Fed. Rep. Ger.
SOURCE: Monatshefte fuer Chemie (1967), 98(4), 1618-50
CODEN: MOCHAP

DOCUMENT TYPE: Journal
LANGUAGE: German
OTHER SOURCE(S): CASREACT 67:116333

GI For diagram(s), see printed CA Issue.

AB cf. CA 67: 99380s. Disubstituted nitrilimines (16), released from hydrazide halides with Et3N were compared in their ability to undergo cycloaddns. with 12 dipolarophiles of various activities. Electron-attracting substituents on the nitrilimine C and N stabilized the ground state, reduced the 1,3-dipolar activity, and promoted the formation of tetrasubstituted 1,4-dihydro-1,2,4,5-tetrazines (I). At least 3

different reaction paths from hydrazide halides to 1,4-dihydrotetrazines were shown.

L5 ANSWER 29 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1966:67089 CAPLUS
DOCUMENT NUMBER: 64:67089
ORIGINAL REFERENCE NO.: 64:12503b-c
TITLE: Pyrazoles. LIII. Ultraviolet and fluorescence spectra of some phenylpyrazoles. Ortho effect of substituents in position 5
AUTHOR(S): Grandberg, I. I.; Tabak, S. V.; Kost, A. N.
CORPORATE SOURCE: M. V. Lomonosov State Univ., Moscow
SOURCE: Khimiya Geterotsiklicheskikh Soedinenii (1965), (6), 901-4
CODEN: KGSSAQ; ISSN: 0132-6244
DOCUMENT TYPE: Journal
LANGUAGE: Russian

AB cf. CA 64, 0704c. Uv absorption spectra in MeOH of 1-phenylpyrazoles substituted in position 5 differed from derivs. substituted in positions 3 and 4 by hypsochromic shifts with hypochromic effect. This fact was explained by the effect of both electron-donor and electron-acceptor substituents in position 5 of pyrazole on the phenyl ring. The coplanarity of the phenyl and pyrazole rings violated the ortho effect. The groups increasing the steric effect were found in the order: Ph < OH equal or nearly equal to NH₂ < Cl < NHAc < Me < CO₂H. Analysis of the fluorescence spectra proved this fact. 19 references.

L5 ANSWER 30 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1964:432381 CAPLUS
DOCUMENT NUMBER: 61:32381
ORIGINAL REFERENCE NO.: 61:5631g-h, 5632a
TITLE: Reactions of phenyl-substituted heterocyclic compounds. V. Nitrations of 1,3- and 1,5-diphenylpyrazoles
AUTHOR(S): Lynch, Brian M.; Hung, Yuk-Yung
CORPORATE SOURCE: St. Francis Xavier Univ., Antigonish
SOURCE: Canadian Journal of Chemistry (1964), 42(7), 1605-15
CODEN: CJCHAG; ISSN: 0008-4042
DOCUMENT TYPE: Journal
LANGUAGE: Unavailable
OTHER SOURCE(S): CASREACT 61:32381

GI For diagram(s), see printed CA Issue.

AB cf. CA 59, 10053c. Dinitration of 1,3- or 1,5-diphenylpyrazole in H₂SO₄ gave the corresponding bis(p-nitrophenyl) compds. (I) and (II), while HNO₃-Ac₂O gave the 4-nitro-1-(p-nitrophenyl)-compds. (III and IV). Mononitration at the 4-position occurred when the diphenylpyrazoles and several other 1-phenyl-pyrazoles were nitrated at 0° by HNO₃-Ac₂O. Possible explanations of the dependence of orientation on the nature of the nitrating agent were discussed. Nuclear magnetic resonance (n.m.r.) spectroscopy was used in demonstrating the structures of many of the nitration products, and a general discussion of the n.m.r. spectra of substituted 1-phenylpyrazoles was given.

L5 ANSWER 31 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1963:408935 CAPLUS
DOCUMENT NUMBER: 59:8935
ORIGINAL REFERENCE NO.: 59:1616a-c
TITLE: Pyrazoles. XXXIV. Ultraviolet spectra of pyrazole systems
AUTHOR(S): Grandberg, I. I.
CORPORATE SOURCE: M. V. Lomonosov State Univ., Moscow
SOURCE: Zhurnal Obshchei Khimii (1963), 33, 519-25

CODEN: ZOKHA4; ISSN: 0044-460X

DOCUMENT TYPE: Journal
LANGUAGE: Unavailable

AB Ultraviolet spectra are reported for 117 substituted pyrazoles. Halogen atoms, alkyl, or NH₂ groups produce a small bathochromic effect on the K band of pyrazole; in the presence of only these auxochromes the band is below 235 mμ; chromophores such as aryl groups, NO₂, or NO groups, caused a shift of the K band to 242-80 mμ. The largest bathochromic shift occurs with auxochromes in 1- and 4-positions. If the group interaction between these substituents and the ring occurs through p-electrons, the bathochromic shift is small. Estimation of electron mobilities of heterocyclic rings on the basis of bathochromic band shifts resulted in the following series of ascending magnitude of the shift: 2-selenophene-yl, 2-thienyl, 2-furyl, Ph. The ferrocenyl radical as a substituent on the pyrazole ring acts as a typical auxochrome and does not conjugate with the pyrazole ring. Cf. CA 58,3290f.

L5 ANSWER 32 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1963:402891 CAPLUS

DOCUMENT NUMBER: 59:2891

ORIGINAL REFERENCE NO.: 59:414e-g

TITLE: Reactions of phenyl-substituted heterocyclic compounds. II. Nitrations and brominations of 1-phenylpyrazole derivatives

AUTHOR(S): Khan, Misbahul Ain; Lynch, Brian M.; Hung, Yuk-Yung

CORPORATE SOURCE: Mem. Univ. Newfoundland, St. John's

SOURCE: Canadian Journal of Chemistry (1963), 41, 1540-7

CODEN: CJCHAG; ISSN: 0008-4042

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

GI For diagram(s), see printed CA Issue.

AB Nitrations of 1-phenylpyrazole (I, R = H) (II) 1-p-biphenylpyrazole (III) and 1,5-diphenylpyrazole by acetyl nitrate (nitric acid-acetic anhydride) occur selectively in the 4-position of the pyrazole ring, as do brominations of II and III in CHCl₃. These results are in agreement with Brown's calcns. (CA 49, 8642g) of localization energies for electrophilic substitution in pyrazole. II, e.g., gives I (R = NO₂). However, nitration of II by mixed acids at 12° yields 1-p-nitrophenylpyrazole, and bromination of II by Br in concentrated H₂SO₄ in the presence of Ag₂SO₄ yields 1-p-bromophenylpyrazole. The variations in orientation of substitution can be rationalized if the reacting species of I in strongly acidic solvents is the conjugate acid, in which the pyrazole ring is deactivated by protonation. Cf. CA 58, 6819f.

L5 ANSWER 33 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1960:2188 CAPLUS

DOCUMENT NUMBER: 54:2188

ORIGINAL REFERENCE NO.: 54:505b-e

TITLE: Pyrazoles. II. Synthesis of N-phenylpyrazoles from corresponding pyrazolines

AUTHOR(S): Grandberg, I. I.; Kost, A. N.

CORPORATE SOURCE: State Univ., Moscow

SOURCE: Zhurnal Obshchei Khimii (1959), 29, 658-62

CODEN: ZOKHA4; ISSN: 0044-460X

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

OTHER SOURCE(S): CASREACT 54:2188

AB cf. C.A. 53, 10188f. Adding over 1.5 hrs. of 58 g. CH₂:CHCHO and 108 g. PhNHNH₂ to 400 ml. Et₂O at 5-10°, stirring 5 hrs., evaporating the solvent, adding 300 ml. 2N H₂SO₄ and steam distilling gave 30% N-phenylpyrazoline, b₁₄ 143-6°, n_{20D} 1.6156, d₂₀ 1.0984. This (29.2 g.) refluxed gently 1 hr. with 6.4 g. S gave 76% 1-phenylpyrazole,

b6 106-9°, b10 124°, b30 141-2°, 1.5976, 1.0908.
 Similarly were obtained: 61% 1-phenyl-3-methylpyrazole, b19 139-40°, m. 38°; 88% 1,3-diphenylpyrazole, m. 85°; 84% 1,5-diphenylpyrazole, b8 182-3°, m. 56°. Heating 0.5 hr. 20.8 g. benzalacetophenone with 10.8 g. PhNHNH2 in 15 ml. EtOH and 15 ml. C6H6 (if an exothermic reaction failed to occur, 2 drops HCO2H was added), the solvents removed and the residue heated 1 hr. with 3.1 g. S as above yielded 67% 1,3,5-triphenylpyrazole, m. 140°. Similarly were obtained: 60% 1-phenyl-3-tert-butylpyrazole, b18 151°, m. 74°; 87% 1-phenyl-3-acetamidopyrazole, m. 128°; and 39% 1-phenyl-3-aminopyrazole, b8 167-70°, m. 90° (picrate, m. 158-9°) (Duffin and Kendall, C.A. 49, 10269c).

L5 ANSWER 34 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1948:25374 CAPLUS

DOCUMENT NUMBER: 42:25374

ORIGINAL REFERENCE NO.: 42:5450f-i, 5451a-i, 5452a-i, 5453a-d

TITLE: Heterocyclic syntheses. IX. Ketone reagents and anils of hydroxymethylene ketones

AUTHOR(S): Panizzi, Luigi; Monti, Elios

CORPORATE SOURCE: Ist. chim. generale anal. politec., Milan, Italy

SOURCE: Gazzetta Chimica Italiana (1947), 77, 556-71

CODEN: GCITA9; ISSN: 0016-5603

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

GI For diagram(s), see printed CA Issue.

AB cf. C.A. 42, 903h. Whereas NH2OH.HCl (I) and PhNHNH2 (II) react with both the CO and the CHOH group of RCOCH:CHOH compds., and form 3- and 5-substituted isoxazoles and pyrazoles, with RCOCH:CHOR' (III) compds. the reaction is confined to the CO group, whereby only 3-substituted heterocyclic derivs. are formed (cf. P. and Sbrillo-Siena, C.A. 41, 1221d). The present work describes a method for preparing exclusively the corresponding 5 substituted derivs., viz., by making I and II react with RCOCH:CHNHPH (IV) compds. In each case PhNH2 (V) and water are evolved, and cyclization then takes place. The IV structure is preferred to the RCOCH2CH:NPh (VI) structure because it is in better accord with the notable stability to heat, acids, and alkalies, with the formation of similar compds. from secondary anilines, and with spectrochem. measurements of analogous imino-enol-amine systems. However, if the compds. react also in the VI form, the mechanism is probably: A comparison of this reaction with that of III compds., in which tautomerism is impossible, indicates that the presence in IV of the N and of a mobile amino-H has a decisive role in the course of the reaction. The problem should be resolved by the behavior of ketone derivs. formed from hydroxymethylene ketones and secondary amines, where again tautomerism would be impossible. HCO2Et (22 g.) and 14.5 g. acetone, added slowly to a suspension of 5.7 g. powdered Na in 120 cc. anhydrous C6H6, allowed to stand several hrs. at 30-40°, agitated with ice-water, the aqueous layer treated with excess V in AcOH, the orange-brown oil which seps. extracted with C6H6, the extract dried by CaCl2, evaporated, and distilled in vacuo, and the fraction (13 g.) which b12 148-50° allowed to solidify, washed with ligroin, and purified by C6H6-ligroin, yields acetylacetaldehyde anil, AcCH:CHNHPH (VII), m. 50-2°; FeCl3 turns its alc. solns. red. Alc.VII (1 g. in 5 cc.) and 0.65 g. I in a min. of water, refluxed 1.5 hrs., diluted with water, acidified (Congo red) with HCl, extracted with Et2O, the extract distilled, the fraction at 50° agitated with saturated aqueous CdCl2, and the addition product washed with EtOH and Et2O, dried, and distilled twice, yield 0.3 g. of 5-methylisoxazole (VIII). VIII (0.24 g.) and EtONa (from 0.1 g. Na and 1.5 cc. anhydrous EtOH), allowed to stand, diluted with water, 0.55 g. p-O2NC6H4NHNH2 (IX) in 3 cc. glacial AcOH added, then NaCl, allowed to stand 1 hr., filtered, and washed with water, yield

p-O₂NC₆H₄NHN:CM₂CH₂CN, m. 183-5° (cf. Justoni, C.A. 35, 5110.8).
 VII (1 g.), 0.71 g. II, 0.65 cc. concentrated HCl, and 10 cc. EtOH, refluxed 3 hrs., diluted with water, acidified (Congo red) with HCl, extracted with Et₂O, the extract evaporated, the residue steam-distilled, the oil distillate extracted with Et₂O, and the extract dried by Na₂SO₄ and distilled, leave a residue of PhN.N:CH.CH:CM₂. Its chloroplatinate m. 193-6° (decomposition) (cf. Ber. 32, 2891(1899); Stoermer, C.A. 1, 1287), and its picrate m. 93-7° (cf. Ber. 32, 2891(1899); Stoermer, loc. cit.). Alc. VII (0.5 g. in 10 cc.), 1 g. NaOAc, and p-O₂NC₆H₄N₂Cl (X) (from 0.5 g. IX), allowed to stand, and the precipitate purified by BuOH, yield (p-nitrophenylazo)acetylacetaldehyde anil, AcC(:CHNHP₂)N:NC₆H₄NO₂-p (XI), orange, m. 186-8° (decomposition); NaOH turns its alc. solns. intense red. An alc. suspension of XI (1 g. in 110 cc.) and 0.26 g. I, refluxed 3 hrs., allowed to stand, and the precipitate purified by BuOH, yield (p-nitrophenylazo)acetylacetaldehyde oxime, p-O₂NC₆H₄N:NCHAcC(:NOH)H, orange-red, m. 221-3° (decomposition). Alc. VII (1 g. in 15 cc.), 1.8 g. NaOAc, and PhN₂Cl (from 0.6 g. V), allowed to stand and the precipitate purified by EtOH, yield (phenylazo)acetylacetaldehyde anil (XII), yellow, m. 128-30°. XII (0.5 g.), 15 cc. glacial AcOH, and 0.2 g. II, heated 1.5 hrs. on a steam bath and allowed to stand, precipitate the hydrazone, AcCH(N:NPh)CH:NNHP₂, golden yellow, m. 215-18°. The mother liquor, diluted, allowed to stand, and the precipitate purified by MeOH, yields PhN.CH:CH.C(N:NPh):CM₂, m. 108-11°. PhAc (24 g.) and 14 g. HCO₂Me, added slowly to a suspension of 5 g. powdered Na in 100 cc. anhydrous C₆H₆ (the reaction is energetic and must be cooled), allowed to stand, ice water added, a small excess of V.AcOH added to the aqueous layer, and the precipitate purified by BuOH, yields 36 g. BzCH:CHNHP₂ (XIII), lemon-yellow, m. 140-1° (cf. Claisen and Fischer, Ber. 21, 1137(1888)). Alc. XIII (4 g. in 20 cc.) and 1.9 g. I in a min. of water, heated 1 hr. at 100°, most of the EtOH evaporated, diluted with water, acidified (Congo red) with HCl, extracted with Et₂O, the extract dried by Na₂SO₄, evaporated, and the residue (2.5 g.) fractionally distilled, yield 5-phenylisoxazole (XIV), b₃-4 110° (cf. Claisen, Ber. 36, 3671(1909)). XIV (0.496 g.) and EtONa (from 0.35 g. Na and 5 cc. anhydrous EtOH), heated a short time at 40-50°, excess IX in AcOH added, allowed to stand overnight, and the precipitate washed and purified by BuOH, yield α-cyanoacetophenone p-nitrophenylhydrazone, p-O₂NC₆H₄NHN:CPhCH₂CN, yellow, m. 177-8°. XIII (3 g.), 1.74 g. II, 1.5 cc. concentrated HCl, and 40 cc. EtOH, refluxed 2 hrs., most of the EtOH evaporated, diluted with water, acidified with HCl (Congo red), extracted with Et₂O, the extract evaporated, and the oil (2.6 g.) purified by distillation in vacuo, yield PhN.N:CH.CH:CPh (cf. Claisen and Fischer, loc. cit.). Reduction by Na and EtOH yields PhN.N:CH.CH₂CHPh, m. 133-5° (cf. Ber. 26, 112(1893)). XIII (3 g. in 300 cc. MeOH), excess NaOAc, and X (from 1.9 g. p-O₂NC₆H₄NH₂), allowed to stand, and the precipitate purified by BuOH, yield (p-nitrophenylazo)benzoylacetaldehyde anil (XV), orange-red, m. 202-3°. XV (0.46 g.) and 0.1 g. I in 50 cc. EtOH, refluxed 1.5 hrs., evaporated, diluted with water, and the precipitate purified by BuOH, yield (p-nitrophenylazo)benzoylacetaldehyde oxime, m. 209-12°. Alkalies turn its alc. solns. orange-red. XIII (1 g.) in 150 cc. MeOH, 1.5 g. NaOAc, and PhN₂Cl (from 0.5 g. V), allowed to stand, and the precipitate (0.3 g.) purified by EtOH, yield (phenylazo)benzoylacetaldehyde anil (XVI), orange-yellow, m. 137-9°. XVI (0.22 g.), 0.07 g. II, and 15 cc. glacial AcOH, heated 2 hrs. at 100°, diluted with water, partially neutralized, and the precipitate (0.17 g.) purified by EtOH, yield 1,5-diphenyl-4-phenylazopyrazole, PhN.CH:CH.C(N:NPh):CPh, yellow, m. 117-18°. PhCH:CHAc (7.3 g.), 4

g. HCO₂Me, and a suspension of 1.25 g. powdered Na in 60 cc. anhydrous C₆H₆ react energetically and the mixture must be cooled; the product, allowed to stand, agitated with ice-water, excess V.ACOH added to the aqueous layer, and the precipitate purified by EtOH, yield approx. 20% of cinnamoylacetaldehyde anil, PhCH:CHCOCH:CHNHPH (XVII), yellow, m. 150-1°. XVII (2 g.) and 0.8 g. I in 25 cc. EtOH, refluxed 2 hrs., concentrated to a small volume, diluted with water, acidified (Congo red) with HCl, extracted with Et₂O, the extract evaporated, the residue steam-distilled, the distillate allowed to solidify, and purified by petr. ether, yield 5-styrylisoxazole, O.N:CH.CH:CCH:CHPh (XVIII), m. 42-3°; its acetone solution decolorizes KMnO₄; its AcOH solution decolorizes Br slowly. By treatment with cold EtONa solution, dilution with water, acidification, and purification by CCl₄, XVIII forms cinnamoylacetonitrile, PhCH:CHCOCH₂CN, m. 95-8°. With excess IX in AcOH, it ppts. the p-nitrophenylhydrazone, m. 210-12° (cf. Musante, C.A. 37, 2737.5). XVIII (2 g.), 0.95 g. II, 0.85 cc. concentrated HCl, and 15 cc. EtOH, refluxed 2 hrs., concentrated to a small volume, diluted with water, extracted with Et₂O, the extract evaporated, the residue distilled in vacuo, and the distillate, b₁₅₋₂₀ 230°, allowed to solidify and purified by EtOH, yield 1-phenyl-5-styrylpyrazole (XIX), m. 127°, soluble in aqueous HCl. (NH₄)₂Cr₂O₇ (1.5 g.), added slowly to 0.8 g. XIX in 20 cc. boiling 20% H₂SO₄, extracted with Et₂O, the extract evaporated, the residue taken up in aqueous Na₂CO₃, extracted with Et₂O, the aqueous residue acidified with HCl, extracted with Et₂O, the extract evaporated, and the residue heated at 110° (to remove BzOH) and purified by boiling water, yields 0.2 g. 1-phenyl-5-pyrazolecarboxylic acid, PhN.N:CH.CH:CCO₂H, m. 179-81°. X (from 0.3 g. p-O₂NC₆H₄NH₂), added to 0.5 g. XVII in 50 cc. MeOH and 1 g. NaOAc, and the precipitate (0.5 g.) purified by BuOH, yields (p-nitrophenylazo)cinnamoylacetaldehyde anil (XX), orange, m. 161-3°. XX (0.4 g.) in 50 cc. EtOH and 0.1 g. I, refluxed 2 hrs., concentrated to a small volume, and the residue allowed to solidify and purified by BuOH, yield (p-nitrophenylazo)cinnamoylacetaldehyde oxime, PhCH:CHCOCH(N:NC₆H₄NO₂-p)C(:NOH)H, yellow, m. 194°. XVII (1 g.), 1.2 g. NaOAc, and PhN₂Cl (from 0.4 g. V) in 100 cc. MeOH, allowed to stand 1 hr., and the precipitate purified by BuOH, yield (phenylazo)cinnamoylacetaldehyde anil (XXI), red, m. 148-9°. Alc. XXI (0.25 g. in 50 cc.), 0.1 g. II, and 0.1 cc. concentrated HCl, boiled a short time, allowed to stand, and the precipitate purified by BuOH, yield the phenylhydrazone, C₂₃H₂₀ON₄, orange-yellow, m. 215-16°. When heated cautiously in vacuo, and the distillate purified by EtOH, it yields 1-phenyl-4-phenylazo-5-styrylpyrazole, yellow, m. 158-60°; a trace turns concentrated H₂SO₄ intense cherry-red. To study IV compds. in which R is Me₂C:CH-, Me₂C:CHCOCH₂CHO (XXII) was made to react with V.ACOH with the intention of obtaining Me₂C:CHCOCH:CHNHPH. However, the reaction was different and an isomer was obtained. Me₂C:CHAc (20 g.), 16 g. HCO₂Me, 100 cc. anhydrous C₆H₆, and MeONa (from 4.9 g. Na), kept below 10° overnight, agitated with ice-water, the aqueous layer treated with V.ACOH, the brown-red oil extracted with C₆H₆, the extract evaporated, the residue distilled in vacuo, the orange-red fraction, which b₁₄ 150-200°, allowed to partially solidify, filtered, and washed with ligroin, and the residue (6.5 g.), purified by CCl₄, yields 1-phenyl-2,3-dehydro-6,6-dimethyl-4-piperidone, HC:CH.CO.CH₂.CMe₂.NPh (XXIII), m. 132°, soluble in dilute HCl (repptd. unaltered by alkalies); its CS₂ solution absorbs Br; it does not immediately decolorize KMnO₄ in acetone. XXIII (0.5 g.) in 5 cc. MeOH and 0.26 g. I, refluxed 3 hrs.,

diluted with water, extracted with Et₂O, the aqueous layer made alkaline with NaOH, and the green-yellow precipitate (0.3 g.) purified by animal charcoal and ligroin, yield the oxime, HC:CH.C(:NOH).CH₂.CMe₂.NPh, m. 167-9°, soluble in dilute HCl (repptd. by alkalies). XXIII (0.5 g.), 1 g. NaOAc, and p-O₂NC₆H₄N₂Cl (from 0.35 g. p-O₂NC₆H₄NH₂) give 0.5 g. of a precipitate which, purified by EtOH, yields the p-nitrophenylazo derivative, HC:CH.CO.CH₂.CMe₂.NC₆H₄N:NC₆H₄NO₂-p (XXIV), carmine-red, m. 170°; its alc. solns. turn orange-red with NaOH. The constitution of XXIV seems, in view of the similarity between XXIII and dialkylanilines, more probable than that of a derivative formed by coupling on the piperidone nucleus.

L5 ANSWER 35 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1945:12004 CAPLUS

DOCUMENT NUMBER: 39:12004

ORIGINAL REFERENCE NO.: 39:1871c-g

TITLE: Heterocyclic syntheses. IV. New isoxazole- and pyrazolecarboxylic acids

AUTHOR(S): Panizzi, Luigi

SOURCE: Gazzetta Chimica Italiana (1943), 73, 13-23

From: Chem. Zentr. 1944, I, 424-5.

CODEN: GCITA9; ISSN: 0016-5603

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

GI For diagram(s), see printed CA Issue.

AB cf. C.A. 38, 5500.2. BzCH₂CO₂Et, HC(OEt)₃ and Ac₂O give Et (ethoxymethylene)benzoylacetate (I), light yellow oil, b₈-10 192-5°; warming with FeCl₃ gives a red color. I with absolute EtOH-KOH in ether at 0° gives Et (hydroxymethylene)benzoylacetate, yellow, b₃-4 135-6° (partial decomposition). I and NH₂OH in EtOH, boiled 2 hrs., give the Et ester (II), b₄₅ 142-3°, of 5-phenyl-4-isoxazolecarboxylic acid (III), O.N:CH.C(CO₂H):CPh, m. 155-6° (prepared from the ester with 1:1 HCl). The structure of III follows from the reaction of II with boiling alc. KOH or with cold EtONa to give Et benzoylcianoacetate, m. 40-1°. I and PhNHNH₂ in AcOH give the Et ester, m. 112.5-14°, of 1,5-diphenyl-4-pyrazolecarboxylic acid (IV), N:CH.C(CO₂H):CPh.NPh, m. 180° (decomposition); decarboxylation of IV yields 1,5-diphenylpyrazole, which was reduced by EtONa to the pyrazoline, m. 135-6°. The positions of the Ph groups in III and IV are taken as proof that NH₂OH and PhNHNH₂ add to the double bond and not to the CO group; ring formation follows through the loss of EtOH and H₂O. Et phenylpyruvate (m. 45°) yields with HC(OEt)₃ and Ac₂O Et (ethoxymethylene)phenylpyruvate, b₃-4 171-2°; the free acid (V) m. 124°. V and NH₂OH in EtOH give a compound (VI) [assumed to be PhCH(COCO₂Et)CH(OEt)NHOH], thick reddish oil, b₃-4 155-8° (partial decomposition); the alkali solubility and FeCl₃ reaction indicate an enol form; with 20% KOH VI gives a small yield of 4-phenyl-5-isoxazolecarboxylic acid, m. 158° (decomposition); the Et ester (an oil) and EtONa give PhCH(CN)COCO₂Et(?), which was not characterized. V and PhNHNH₂ yield an amorphous product which with alkali gives 1,4-diphenyl-3-pyrazolecarboxylic acid, m. 227-8°.

L5 ANSWER 36 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1941:17981 CAPLUS

DOCUMENT NUMBER: 35:17981

ORIGINAL REFERENCE NO.: 35:2891b-f

TITLE: Transformation of 3-isoxazolecarboxylic acids into pyrazole derivatives. IV

AUTHOR(S): Cusmano, Sigismondo

SOURCE: Gazzetta Chimica Italiana (1940), 70, 227-35
CODEN: GCITA9; ISSN: 0016-5603

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB cf. C. A. 34, 7903.8. The transformation of 3-isoxazolecarboxylic acids into pyrazolonimines by fusion with PhNHNH₂ may proceed by decarboxylation followed by ring closure of the resulting cyano ketone phenylhydrazone. To test this hypothesis the fusion was repeated in the presence of Natur Kupfer C (I) (or ordinary reduced Cu) so that, at the lower decarboxylation temps. it might be possible to isolate the phenylhydrazone prior to ring closure and so shed some light on the mechanism of the reaction. A mixture of 1 g. of 5-phenyl-3-isoxazolecarboxylic acid (II), 1 g. I and 1 g. PhNHNH₂ in 20 cc. alc. was boiled for a few min. over a free flame, filtered, alkalinized with Na₂CO₃, extracted free from PhNHNH₂ with ether, acidified with dilute H₂SO₄, and extracted with ether. The residue from the evaporated extract gave 1,5-diphenyl-3-pyrazolecarboxylic acid (III), m. 185° (Et ester, m. 98°), decarboxylated by fusion to give 1,5-diphenylpyrazole, m. 55°, and identical with the known acid prepared by the action of PhNHNH₂ on BzCH₂COCO₂H. A similar transformation of 5-methyl-3-isoxazolecarboxylic acid (IV) gave 1-phenyl-5-methyl-3-pyrazolecarboxylic acid, m. 136° (Me ester, m. 55°), decarboxylated to 1-phenyl-5-methylpyrazole, transformed into the known picrate, m. 98°. In these transformations alc. can be replaced by other solvents. In the absence of I or in the presence of PhNH₂ instead of PhNHNH₂ the isoxazolecarboxylic acid is recovered unchanged.

L5 ANSWER 37 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1927:11948 CAPLUS

DOCUMENT NUMBER: 21:11948

ORIGINAL REFERENCE NO.: 21:1450c-h

TITLE: Reaction of hydrazine with hydroxymethylene ketones and their derivatives

AUTHOR(S): v. Auwers, K.; Mauss, H.

SOURCE: Justus Liebigs Annalen der Chemie (1927), 452, 182-210
CODEN: JLACBF; ISSN: 0075-4617

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB BzCH:CHONa(I) and PhNHNH₂ give a crude product, m. 110-2°, which, recrystd. from C₆H₆ and EtOH, gives 2-[β-benzoylvinyl]-1-phenylhydrazine (II), m. 129.5-30.5°; heating at 185-7° and 14 mm. gives 1,5-diphenylpyrazole (III), m. 55-6°; the mother liquor gives a mixture of III and the 1,3-di-Ph derivative (IV). Attempts to benzoylate II gives only III. II does not react with I; it is oxidized by FeCl₃ but the only product isolated was III; II in EtOH-NaOH, shaken with air, gives β-[benzoylvinyl]azobenzene, red, m. 50-60°. I and PhNHNH₂ in EtOH at room temperature for 2-3 days give PhNHNHCHO (V), m. 141.5-25°, pale yellow crystals, m. 146-7°; II, the azo derivative and BzMe. Crystallizing II from Me₂CO, AcOEt or C₆H₆ gives the isomeric 1-[β-benzoylvinyl]-1-phenylhydrazine (V), m. 164° while mother liquor gives IV. II or V and p-O₂NC₆H₄NHNH₂ in C₅H₅N for 2 hrs. at room temperature give the compound BzCH:CHNPhNHCOC₆H₄NO₂, yellow, m. 205°. Oxidation of II or VI with KMnO₄ in Me₂CO gives BzCH:CHNPh. Condensation of II or VI with p-O₂NC₆H₄CHO gives the p-nitrobenzal derivative of VI, yellow, m. 173-4°, also obtained from BzCH:CHOH and p-O₂NC₆H₄CH:NNHPh. BzCH:CHOH and V in EtOH, heated 2 hrs., give the α-phenyl-β-formylhydrazide of hydroxymethylacetophenone, yellow, m. 148-9°; heated with 2 N HCl it yields IV. I and p-O₂NC₆H₄NHNH₂.HCl in dilute EtOH give the p-nitrophenylhydrazide, orange-yellow, m. 156-7°; heating with AcOH about 0.25 hr. gives 1-[p-Nitrophenyl]-5-phenylpyrazole (VII), b₁₄ 240-2°, m. 117-8°; this also results from p-O₂NC₆H₄NHNH₂ and

BzCH:CHOBz. The p-amino derivative, m. 148.5-9.5°; the diazo compound treated with SnCl₂ and NaOH, gives III. When BzCH:CHOBz and p-O₂NC₆H₄NHNH₂ in PrOH are allowed to stand 1 day there results some VII and the bis-p-nitrophenylhydrazine derivative, orange-red, m. 195-200°, crystallizing with 1 Me₂CO; heating on the H₂O bath gives 1-[p-nitrophenyl]-3-phenylpyrazole (VIII), yellow, m. 169-9.5°. 1-[p-Nitrophenyl]-3-phenylpyrazol-5-one, brownish, m. 202-3°; attempted reduction with PBr₃, resulted in complete decomposition Nitration of VIII or IV gives the same dinitro derivative, m. 226-8°. BzCH:CHNHNMeCH:CHBz, m. 137.5-8.5°, and Me₂SO₄ give a di-Me derivative, m. 175-6°, decomposed by heating with mineral acid or alkali; the neutral product is C₆H₃BZ₃, m. 119-20°. The di-Me derivative was obtained from I and (NHMe)₂.2HCl in HCl. 1-Methyl-1-[benzoylvinyl]-2-[p-nitrobenzal]hydrazine, yellow, m. 207°.

L5 ANSWER 38 OF 38 CAPLUS COPYRIGHT 2009 ACS on STN

ACCESSION NUMBER: 1925:15752 CAPLUS
 DOCUMENT NUMBER: 19:15752
 ORIGINAL REFERENCE NO.: 19:2048f-i,2049a-g
 TITLE: Isomerism relationships in the pyrazole series. II. 3(5)-Phenylpyrazole and its derivatives
 AUTHOR(S): v. Auwers, K.; Schmidt, W.
 SOURCE: Berichte der Deutschen Chemischen Gesellschaft [Abteilung] B: Abhandlungen (1925), 58B, 528-43
 CODEN: BDCBAD; ISSN: 0365-9488
 DOCUMENT TYPE: Journal
 LANGUAGE: Unavailable
 OTHER SOURCE(S): CASREACT 19:15752

GI For diagram(s), see printed CA Issue.

AB cf. C. A. 18, 1296. Unlike the dialkylpyrazoles, of which the 1,5-forms are incapable of existence, the C-phenylalkylpyrazoles can exist in both the 1,3- and 1,5-forms. While it might be concluded from this that the non-existence of the 1,5-dialkyl derivs. is due, as had been tentatively suggested, to a mutual repulsion of the electrochem. similar radicals, the relationships are not so simple as all this and further data will be necessary before the phenomena can be explained. Both 1,3- (I) and 1,5-diphenylpyrazole (II) are stable and their structures have been determined with certainty, but Claisen and Fischer from HOCH:CHCOPh (III) and Ph₂ in Et₂O obtained a compound m. 118-20° which on distillation smoothly yielded II (Ber. 21, 1139(1888)) while Knorr and Duden obtained only I (Ber. 27, 109(1893)). v. A. and S. in their own expts. regularly obtained a substance (IV), m. 126°, having the composition of a phenylhydrazone of BzCH₂CHO and giving II on distillation in vacuo. The yield of this product, however, was only 25% and the Et₂O mother liquor on standing a long time deposited yellowish needles, m. 100-3°, which on distillation gave a semi-solid mass from which I was obtained by steam distillation. In the action of PhNHNH₂ on III, therefore, there are formed both of the possible hydrazones BzCH₂CH:NNHPh (IV) and PhC(:NNHPh)CH₂CHO, which explains the apparent discrepancy between the results of C. and F. and of K. and D. BzCH:CHOBz (V), BzCH:CHOCO₂Et (VI) or BzCH:CHOEt (VII) treated with PhNHNH₂ regularly lost the radical attached to the O, even under the mildest conditions, yielding a product C₁₆H₁₄ON₂ (VIII), m. 162°, which is readily converted into I but does not give PhNH₂ with Na-Hg and AcOH in alc., is insol. in alkalies and does not show aldehyde reactions; it is provisionally assigned the structure PhN.N:CPh.CH₂.CHOH. When it is treated in C₅H₅N with ClCO₂Et a H atom is replaced by CO₂Et and the resulting ester (IX) is readily hydrolyzed. In preparing VIII, the crude product frequently m. around 115-20° but a single crystallization sufficed to raise the m. p. to 162°; v. A. and S. think it likely that the low melting product is a labile form of VIII. VIII smoothly yields I when warmed a very short time with glacial AcOH or kept 5 min. in it in the

cold or when warmed a short time with alc. and a couple drops of HCl or when boiled in Me₂CO, while IV is converted into II only on long boiling in glacial AcOH or dilute alc. HCl or on distillation, indicating that of the 2 isomers I has the greater tendency to be formed. 3(5)-Phenylpyrazole (X) with MeBr in a sealed tube or with Me₂SO₄ gives a mixture of chiefly 1-methyl-3-phenylpyrazole (XI) and the 1,5-isomer (XII). XII was also prepared (1) quant. by condensing PhCH:CHBrCHO with MeNHNH₂ and treating the resulting yellow oily bromopyrazoline with NaOAc, and (2) from the Na salt of III, (MeNHNH₂)₂. H₂SO₄ and NaOAc in H₂O or free HI and MeNHNH₂ in Et₂O and treatment of the resulting compound (XIII), (CH:CHBz)₂NNHMe, with hot glacial AcOH or boiling dilute alc. HCl; in the last case there is formed a mixture of XI and XII in which XII predominates. V with MeNHNH₂ gives chiefly XI but also some XII. While according to Wenglein (Diss. Jena, 1895) the methiodide (XIV) obtained from either XI or XII gives only XI on heating, v. A. and S. find that in fact it gives only 32% XI and 68% XII. Ethylation of X with a large excess of EtBr gives only 1 compound (XV), the comparison of whose consts. with those of XI and XII shows clearly that it is 1-ethyl-3-phenylpyrazole. The Na salt of III is obtained in 90% yield (crude product of varying degrees of purity) from 11.5 g. Na wire under cold C₆H₆ slowly treated with 60 g. MeCOPh and 55 g. acid-free HCO₂Et. Bz derivative (V) of III (70% from III and BzCl in C₅H₆N), m. 75-5.5°, unchanged by glacial AcOH or alc. HCl. Et β-benzoylvinyll carbonate (VI) (80% from III and ClCO₂Et in cold C₅H₅N), m. 57-9°. Et β-benzoylvinyll ether (VII), b₁₀ 162-3°. Carboxyethyl derivative (IX) of VIII, m. 156°, rapidly hydrolyzed by dilute H₂SO₄ even in ice. X, b₁₁ 177-8°, b. 313-4°, m. 79°, d_{499.6} 1.0818, n 1.58166, 1.58890, 1.60734 for α, He and β at 100.8°, EΣ 1.09, 1.15, 31% for α, D and β-α. Treated in cold dilute H₂SO₄ with NaNO₂ it gives a yellow oil whose Et₂O solution soon deposits the nitrate of X, m. 126° (decomposition). N-Ac derivative, b₁₀ 157-8°, m. 64-5°. Methiodide of I, m. 172°, regenerates I when heated above its m. p. Methiodide of II, m. 207°, yields II above its m. p. XI m. 55-6°, b₁₂ 145-6°, d_{499.6} 1.0232, n 1.55539, 1.56216, 1.57982 for α, He and β at 100.8°, EΣ 1.27, 1.33, 37% for α, D and α-β. XII b₁₀ 127°, d_{416.8} 1.0946, n 1.58350, 1.58999, 1.60653, 1.62148 for α, He, β and γ at 15.8°, EΣ 0.40, 0.42, 19%, 19% for α, D, β-α and γ-α. Picrate, greenish yellow, m. 143-4°. α-Methyl-β, β-bis-[β'-benzoylvinyll]hydrazine (XIII), yellow, m. 137.5-8.5°, insol. in dilute HCl, soluble (but with rapid alteration) in concentrated acid with golden yellow color. 1-Methyl-3-phenyl-4-bromopyrazole (90% from XI and Br in AcOH), b₁₂ 175-6°, d_{418.9} 1.4707, n 1.61025, 1.61711, 1.63442, 1.65050 for α, He, β and γ at 18.9°, EΣ 0.27, 0.29, 20%, 22% for α, D, β-α and γ-α. 1-Methyl-5-phenyl-4-bromopyrazole b₁₂ 155-6°, m. 63-4° and if quickly cooled if m. 65-6° but if cooled slowly it again m. 53-4°, d_{456.6} 1.4199, n 1.58179, 1.58790, 1.60325, 1.61706 at 57.6°, EΣ 0.20, 0.22, 13%, 13%. Methylation of the 4-Br derivative of X with alkaline Me₂SO₄ gives a mixture of the 2 above compds.

XV b₁₂

152-3°, m. 36.5-7.5°, d_{411.6} 1.0667, n 1.58209, 1.58897, 1.60657, 1.62283 at 11.6°, EΣ 0.92, 0.98, 34%, 36%.

=>